

## Clinical Policy: Inclisiran (Leqvio)

Reference Number: CP.PHAR.568

Effective Date: 03.01.22

Last Review Date: 02.25

Line of Business: Commercial, HIM, Medicaid

[Coding Implications](#)  
[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### Description

Inclisiran (Leqvio<sup>®</sup>) is a small interfering ribonucleic acid (siRNA) directed to proprotein convertase subtilisin kexin type 9 (PCSK9) messenger RNA (mRNA).

### FDA Approved Indication(s)

Leqvio is indicated as an adjunct to diet and exercise to reduce low-density lipoprotein cholesterol (LDL-C) in adults with hypercholesterolemia, including heterozygous familial hypercholesterolemia (HeFH).

### Policy/Criteria

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Leqvio is **medically necessary** when the following criteria are met:

#### I. Initial Approval Criteria

##### A. Primary Hypercholesterolemia (including HeFH) and Atherosclerotic Cardiovascular Disease (must meet all):

1. Diagnosis of one of the following (a, b, or c):
  - a. **HeFH**, and both of the following (i and ii):
    - i. Baseline LDL-C (prior to any lipid-lowering pharmacologic therapy) was one of the following (1 or 2):
      - 1) If age < 20 years:  $\geq 160$  mg/dL;
      - 2) If age  $\geq 20$  years:  $\geq 190$  mg/dL;
    - ii. HeFH diagnosis is confirmed by one of the following (1 or 2):
      - 1) World Health Organization (WHO)/Dutch Lipid Network familial hypercholesterolemia diagnostic criteria score of  $> 8$  as determined by requesting provider (*see Appendix D*);
      - 2) Definite diagnosis per Simon Broome criteria (*see Appendix D*);
  - b. **Primary hypercholesterolemia that is not HeFH**, and both of the following (i and ii):
    - i. Documentation of one of the following (1 or 2):
      - 1) Presence of a genetically mediated form of primary hypercholesterolemia as evidenced by confirmatory genetic testing results;

- 2) A diagnosis of secondary hypercholesterolemia has been ruled out with absence of all of the following potential causes of elevated cholesterol (a-f):
  - a) Poor diet;
  - b) Hypothyroidism;
  - c) Obstructive liver disease;
  - d) Renal liver disease;
  - e) Nephrosis;
  - f) Medications that have had a clinically relevant contributory effect on the current degree of the member's elevated lipid levels including, but not limited to: glucocorticoids, sex hormones, antipsychotics, antiretrovirals, immunosuppressive agents, retinoic acid derivatives;
- ii. Baseline LDL-C (prior to any lipid-lowering pharmacologic therapy) was  $\geq$  190 mg/dL;
- c. **Atherosclerotic cardiovascular disease (ASCVD)** as evidenced by a history of any one of the following conditions (i-vii):
  - i. Acute coronary syndromes;
  - ii. Clinically significant coronary heart disease (CHD) diagnosed by invasive or noninvasive testing (such as coronary angiography, stress test using treadmill, stress echocardiography, or nuclear imaging);
  - iii. Coronary or other arterial revascularization;
  - iv. Myocardial infarction;
  - v. Peripheral arterial disease presumed to be of atherosclerotic origin;
  - vi. Stable or unstable angina;
  - vii. Stroke or transient ischemic attack (TIA);
2. Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist;
3. Age  $\geq$  18 years;
4. Failure of an 8 week trial of a preferred PCSK9 inhibitor, if applicable, at up to maximally indicated doses, unless contraindicated or clinically significant adverse effects are experienced;  
*\*Prior authorization may be required for PCSK9 inhibitors*
5. For members on statin therapy, both of the following (a and b):
  - a. Leqvio is prescribed in conjunction with a statin at the maximally tolerated dose;
  - b. Member has been adherent for at least the last 4 months to maximally tolerated doses of one of the following statin regimens (i, ii, or iii):
    - i. A high intensity statin (*see Appendix E*);
    - ii. A moderate intensity statin (*see Appendix E*), and member has one of the following (1 or 2):
      - 1) Intolerance to two high intensity statins;
      - 2) A statin risk factor (*see Appendix G*);
    - iii. A low intensity statin and member has one of the following (1 or 2):
      - 1) Intolerance to one high and one moderate intensity statins;
      - 2) A statin risk factor (*see Appendix G*) and history of intolerance to two moderate intensity statins;

6. For members not on statin therapy, member meets one of the following (a or b):
  - a. Statin therapy is contraindicated per Appendix F;
  - b. For members who are statin intolerant, both of the following (i and ii):
    - i. Member has tried at least two statins, one of which must be hydrophilic (pravastatin, fluvastatin, or rosuvastatin);
    - ii. Member meets one of the following (1 or 2):
      - 1) Member has documented statin risk factors (*see Appendix G*);
      - 2) Member is statin intolerant due to statin-associated muscle symptoms (SAMS) and meets both of the following (a and b):
        - a) Documentation of intolerable SAMS persisting at least two weeks, which disappeared with discontinuing the statin therapy and recurred with a statin re-challenge;
        - b) Documentation of re-challenge with titration from lowest possible dose and/or intermittent dosing frequency (e.g., 1 to 3 times weekly);
7. Member has been adherent to ezetimibe therapy used concomitantly with a statin at the maximally tolerated dose for at least the last 4 months, unless contraindicated per Appendix F or member has a history of ezetimibe intolerance (e.g., associated diarrhea or upper respiratory tract infection);
8. Documentation of recent (within the last 60 days) LDL-C of one of the following (a or b):
  - a. If member has ASCVD (i or ii):
    - i.  $\geq 70$  mg/dL;
    - ii.  $\geq 55$  mg/dL, and member is at very high risk (*see Appendix I*);
  - b. If member has severe primary hypercholesterolemia (including HeFH):  $\geq 100$  mg/dL;
9. Treatment plan does not include coadministration with Juxtapid<sup>®</sup>, Repatha<sup>®</sup>, or Praluent<sup>®</sup>;
10. Dose does not exceed 284 mg initially and at 3 months, then every 6 months thereafter.

**Approval duration:**

**Medicaid/HIM** – 9 months

**Commercial** – 6 months or to the member’s renewal date, whichever is longer

**B. Other diagnoses/indications (must meet 1 or 2):**

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or

2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

## II. Continued Therapy

### A. Primary Hypercholesterolemia (including HeFH) and Atherosclerotic Cardiovascular Disease (must meet all):

1. Currently meets one of the following (a or b):
  - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
  - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (refer to state specific addendums for *CC.PHARM.03A* and *CC.PHARM.03B*);
2. If statin tolerant, documentation of adherence to a statin at the maximally tolerated dose;
3. Member is responding positively to therapy as evidenced by lab results within the last 3 months showing an LDL-C reduction since initiation of Leqvio therapy;
4. Treatment plan does not include coadministration with Juxtapid, Repatha, or Praluent;
5. If request is for a dose increase, new dose does not exceed 284 mg every 6 months.

#### Approval duration:

**Medicaid/HIM** – 12 months

**Commercial** – 6 months or to the member’s renewal date, whichever is longer

### B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
  - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
  - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

## III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies –

CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

ASCVD: atherosclerotic cardiovascular disease	mRNA: messenger RNA
CHD: coronary heart disease	PCSK9: proprotein convertase subtilisin–kexin type 9
FDA: Food and Drug Administration	RNA: ribonucleic acid
FH: familial hypercholesterolemia	SAMS: statin-associated muscle symptoms
HeFH: heterozygous familial hypercholesterolemia	siRNA: small interfering RNA
LDL-C: low density lipoprotein cholesterol	TIA: transient ischemic attack
	WHO: World Health Organization

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

<b>Drug Name</b>	<b>Dosing Regimen</b>	<b>Dose Limit/ Maximum Dose</b>
ezetimibe/simvastatin (Vytorin <sup>®</sup> )	10/40 mg PO QD	10 mg-40 mg/day (Use of the 10/80 mg dose is restricted to patients who have been taking simvastatin 80 mg for 12 months or more without evidence of muscle toxicity)
ezetimibe (Zetia <sup>®</sup> )	10 mg PO QD	10 mg/day
atorvastatin (Lipitor <sup>®</sup> )	40 mg PO QD	80 mg/day
rosuvastatin (Crestor <sup>®</sup> )	5 - 40 mg PO QD	40 mg/day
Praluent (alirocumab)	75 mg SC once every 2 weeks or 300 mg SC once every 4 weeks; if response to 75 mg every 2 weeks or 300 mg every 4 weeks is inadequate, dose may be increased to 150 mg once every 2 weeks	300 mg/month
Repatha (evolocumab)	140 mg SC every 2 weeks or 420 mg SC once monthly	420 mg/month

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

*Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s): history of serious hypersensitivity reaction to inclisiran or any of the excipients of Leqvio
- Boxed warning(s): none reported

Appendix D: Criteria for Diagnosis of HeFH

- Dutch Lipid Clinic Network criteria for Familial Hypercholesterolemia (FH)

FH Criteria	Points	Member's Score†
<b>Family History</b>		
First-degree relative with known premature* coronary and vascular disease	1	Place highest score here (0, 1 or 2)
First-degree relative with known LDL-C level above the 95 <sup>th</sup> percentile	1	
First-degree relative with tendinous xanthomata and/or arcus cornealis	2	
Children aged < 18 years with LDL-C level above the 95 <sup>th</sup> percentile	2	
<b>Clinical History</b>		
Patient with premature* coronary artery disease	2	Place highest score here (0, 1 or 2)
Patient with premature* cerebral or peripheral vascular disease	1	
<b>Physical Examination</b>		
Tendinous xanthomata	6	Place highest score here (0, 4 or 6)
Arcus cornealis prior to age 45 years	4	
<b>Cholesterol Levels - mg/dL (mmol/liter)</b>		
LDL-C ≥ 330 mg/dL (≥ 8.5)	8	Place highest score here (0, 1, 3, 5 or 8)
LDL-C 250 – 329 mg/dL (6.5 – 8.4)	5	
LDL-C 190 – 249 mg/dL (5.0 – 6.4)	3	
LDL-C 155 – 189 mg/dL (4.0 – 4.9)	1	
<b>DNA Analysis</b>		
Functional mutation in the <i>LDLR</i> , <i>apo B</i> or <i>PCSK9</i> gene	8	Place score here (0 or 8)
<b>TOTAL SCORE</b>	Definite FH: > 8	Place total score here ___

\*Premature – men < 55 years or women < 60 years

†Choose the highest score from each of the five categories and then add together for a total score. The five categories are 1) Family History, 2) Clinical History, 3) Physical Examination, 4) Cholesterol Levels, and 5) DNA Analysis.

- Simon Broome Register Group Definition of Definite FH (meets 1 and 2):
  - One of the following (a or b):
    - Total cholesterol level above 7.5 mmol/L (290 mg/dL) in adults or a total cholesterol level above 6.7 mmol/L (260 mg/dL) for children under 16
    - LDL levels above 4.9 mmol/L (190 mg/dL) in adults (4.0 mmol/l in children) (either pre-treatment or highest on treatment)
  - One of the following (a or b):
    - Tendinous xanthomas in patient or relative (parent, child, sibling, grandparent, aunt, uncle)

- b. DNA-based evidence of an LDL receptor mutation or familial defective apo B-100
- High and Moderate Risk of ASCVD:
  - Patients with high risk of ASCVD include the following:
    - History of clinical atherosclerotic cardiovascular disease (as defined in section II)
    - Diabetes with an estimated 10-year ASCVD risk  $\geq 7.5\%$  for adults 40-75 years of age
    - Untreated LDL  $\geq 190$  mg/dL
  - Patients with moderate risk of ASCVD include the following:
    - Diabetes with an estimated 10-year ASCVD risk  $< 7.5\%$  for adults 40-75 years of age
    - Estimated 10-year ASCVD risk  $\geq 5\%$  for adults 40-75 years of age
  - The calculator for the 10-year ASCVD risk estimator can be found here: <http://tools.acc.org/ASCVD-Risk-Estimator-Plus/#!/calculate/estimate>. Information needed to complete the ASCVD Risk Estimator include: gender, race (white, African American, other), systolic blood pressure, history of diabetes, age, total cholesterol, HDL-cholesterol, treatment for hypertension, smoking history or status, and concurrent statin or aspirin therapy.

*Appendix E: High, Moderate, and Low Intensity Daily Statin Therapy for Adults*

<b>High Intensity Statin Therapy</b> <i>Daily dose shown to lower LDL-C, on average, by approximately <math>\geq 50\%</math></i>
<ul style="list-style-type: none"> <li>• Atorvastatin 40-80 mg</li> <li>• Rosuvastatin 20-40 mg</li> </ul>
<b>Moderate Intensity Statin Therapy</b> <i>Daily dose shown to lower LDL-C, on average, by approximately 30% to 50%</i>
<ul style="list-style-type: none"> <li>• Atorvastatin 10-20 mg</li> <li>• Fluvastatin XL 80 mg</li> <li>• Fluvastatin 40 mg BID</li> <li>• Lovastatin 40 mg</li> <li>• Pitavastatin 1-4 mg</li> <li>• Pravastatin 40-80 mg</li> <li>• Rosuvastatin 5-10 mg</li> <li>• Simvastatin 20-40 mg</li> </ul>
<b>Low Intensity Statin Therapy</b> <i>Daily dose shown to lower LDL-C, on average, by <math>&lt; 30\%</math></i>
<ul style="list-style-type: none"> <li>• Simvastatin 10 mg</li> <li>• Pravastatin 10-20 mg</li> <li>• Lovastatin 20 mg</li> <li>• Fluvastatin 20-40 mg</li> </ul>

*Appendix F: Statin and Ezetimibe Contraindications*

<b>Statins</b>
<ul style="list-style-type: none"> <li>• Decompensated liver disease (development of jaundice, ascites, variceal bleeding, encephalopathy)</li> </ul>

Statins
<ul style="list-style-type: none"> <li>• Laboratory-confirmed acute liver injury or rhabdomyolysis resulting from statin treatment</li> <li>• Pregnancy*, actively trying to become pregnant, or nursing</li> <li>• Immune-mediated hypersensitivity to the HMG-CoA reductase inhibitor drug class (statins) as evidenced by an allergic reaction occurring with at least TWO different statins</li> </ul>
Ezetimibe
<ul style="list-style-type: none"> <li>• Moderate or severe hepatic impairment [Child-Pugh classes B and C]</li> <li>• Hypersensitivity to ezetimibe (e.g., anaphylaxis, angioedema, rash, urticaria)</li> </ul>

*\*In July 2021, the FDA requested removal of the contraindication against use of statins in pregnant women. Because the benefits of statins may include prevention of serious or potentially fatal events in a small group of very high-risk pregnant patients, contraindicating these drugs in all pregnant women is not appropriate. <https://www.fda.gov/safety/medical-product-safety-information/statins-drug-safety-communication-fda-requests-removal-strongest-warning-against-using-cholesterol>*

#### *Appendix G: Statin Risk Factors*

Statin Risk Factors
<ul style="list-style-type: none"> <li>• Multiple or serious comorbidities, including impaired renal or hepatic function</li> <li>• Unexplained alanine transaminase (ALT) elevations &gt; 3 times upper limit of normal, or active liver disease</li> <li>• Concomitant use of drugs adversely affecting statin metabolism</li> <li>• Age &gt; 75 years, or history of hemorrhagic stroke</li> <li>• Asian ancestry</li> </ul>

#### *Appendix H: General Information*

- Patients should remain on concomitant therapy with a statin if tolerated due to the established long term cardiovascular benefits.
- The diagnosis of SAMS is often on the basis of clinical criteria. Typical SAMS include muscle pain and aching (myalgia), cramps, and weakness. Symptoms are usually bilateral and involve large muscle groups, including the thigh, buttock, back, and shoulder girdle musculature. In contrast, cramping is usually unilateral and may involve small muscles of the hands and feet. Symptoms may be more frequent in physically active patients. Symptoms often appear early after starting statin therapy or after an increase in dose and usually resolve or start to dissipate within weeks after cessation of therapy, although it may take several months for symptoms to totally resolve. Persistence of symptoms for more than 2 months after drug cessation should prompt a search for other causes or for underlying muscle disease possibly provoked by statin therapy. The reappearance of symptoms with statin rechallenge and their disappearance with drug cessation offers the best evidence that the symptoms are truly SAMS.
- Pravastatin, fluvastatin, and rosuvastatin are hydrophilic statins which have been reported to confer fewer adverse drug reactions than lipophilic statins.
- In a final evidence report published March 2021, the Institute for Clinical and Economic Review (ICER) concluded that while uncertainty remains regarding the magnitude of overall benefit and how inclisiran compares to that of PCSK9 inhibitors, the current evidence offers high certainty of at least a small net health benefit for inclisiran when

used for patients who have need of significant reduction in LDL-C despite maximally tolerated oral lipid-lowering therapy (B+).

*Appendix I: Criteria for Defining Patients at Very High Risk of Future ASCVD Events<sup>5</sup>*

Very high risk is defined as having either a history of multiple major ASCVD events **OR** 1 major ASCVD event and multiple high-risk conditions:

- Major ASCVD events:
  - Recent acute coronary syndrome (within the past 12 months)
  - History of myocardial infarction (other than recent acute coronary syndrome event listed above)
  - History of ischemic stroke
  - Symptomatic peripheral artery disease (history of claudication with ankle-brachial index < 0.85 or previous revascularization or amputation)
- High-risk conditions:
  - Age ≥ 65 years
  - HeFH
  - History of prior coronary artery bypass surgery or percutaneous coronary intervention outside of the major ASCVD event(s)
  - Diabetes
  - Hypertension
  - Chronic kidney disease (estimated glomerular filtration rate [eGFR] 15-59 mL/min/1.73 m<sup>2</sup>)
  - Current tobacco smoking
  - Persistently elevated LDL-C (LDL-C ≥ 100 mg/dL [≥ 2.6 mmol/L]) despite maximally tolerated statin therapy and ezetimibe
  - History of congestive heart failure

**V. Dosage and Administration**

Indication	Dosing Regimen	Maximum Dose
Hypercholesterolemia (including HeFH)	284 mg SC on initially and at 3 months, then every 6 months thereafter. If a planned dose is missed by more than 3 months, restart with a new dosing schedule. Leqvio should be administered by a healthcare professional	See regimen

**VI. Product Availability**

Single-dose prefilled syringe: 284 mg/1.5 mL (189 mg/mL)

**VII. References**

1. Leqvio Prescribing Information. East Hanover, NJ: Novartis; July 2025. Available at: [www.leqvio.com](http://www.leqvio.com). Accessed August 6, 2025.
2. Raal FJ, Kallen D, Ray KK, et al. Inclisiran for the treatment of heterozygous familial hypercholesterolemia. *N Engl J Med.* 2020; 382: 1520-1530.
3. Ray KK, Wright RS, Kallend D, et al. Two phase 3 trials of inclisiran in patients with elevated LDL cholesterol. *N Engl J Med.* 2020; 382: 1507-1519.

4. Institute for Clinical and Economic Review. Bempedoic acid and inclisiran for patients with heterozygous familial hypercholesterolemia and for secondary prevention of ASCVD: Effectiveness and value. Final evidence report published March 2, 2021. Available at: <https://icer.org/assessment/high-cholesterol-2021>. Accessed November 6, 2024.  
Guidelines
5. Grundy SM, Stone NJ, Bailey AL, et al. 2018 ACC/AHA/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: A report of the American College of Cardiology Foundation/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2018;Nov 10:[Epub ahead of print].
6. Lloyd-Jones DM, Morris PB, Minissian MB, et al. 2017 Focused update of the 2016 ACC expert consensus decision pathway on the role of non-statin therapies for LDL-cholesterol lowering in the management of atherosclerotic cardiovascular disease risk. *J Am Coll Cardiol* 2017; 70(14):1785-1822. <http://dx.doi.org/10.1016/j.jacc.2017.07.745>.
7. Lloyd-Jones DM, Morris PB, Ballintyne CM, et al. 2022 ACC expert consensus decision pathway on the role of nonstatin therapies for LDL-cholesterol lowering in the management of atherosclerotic cardiovascular disease risk: A report of the American College of Cardiology Solution Set Oversight Committee. *J Am Coll Cardiol*. 2022; 80 (14): 1366-1418.
8. Jacobson TA, et al. National Lipid Association recommendations for patient-centered management of dyslipidemia: part 1 – full report. *Journal of Clinical Lipidology*. March-April 2015; 9(2): 129-169. <http://dx.doi.org/10.1016/j.jacl.2015.02.003>.
9. Goldberg AC, Hopkins PN, Toth PP, et al. Familial hypercholesterolemia: screening, diagnosis and management of pediatric and adult patients: clinical guidance from the National Lipid Association Expert Panel on Familial Hypercholesterolemia. *Journal of Clinical Lipidology*. June 2011; 5(3S): 1-15.
10. Al-Rasadi K, Al-Waili K, Al-Sabti HA, et al. Criteria for diagnosis of familial hypercholesterolemia: A comprehensive analysis of the different guidelines, appraising their suitability in the Omani Arab population. *Oman Medical Journal*. 2014; 29(2): 85–91. <http://doi.org/10.5001/omj.2014.22>.
11. Clinical Lipidology Resource Center, sponsored by the National Lipid Association and the Journal of Clinical Lipidology. Genetic classification of dyslipidemia. Available at: <http://nlaresourcecenter.lipidjournal.com/Content/PDFs/Tables/1.pdf>. Accessed November 6, 2024.
12. Virani SS, Newby LK, Arnold SV, et al. 2023 AHA/ACC/ACCP/ASPC/NLA/PCNA Guideline for the Management of Patients With Chronic Coronary Disease: A Report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Practice Guidelines. *Circulation*. 2023 Aug 29;148(9):e9-e119.
13. Rao SV, O'Donoghue ML, Ruel M, et al. 2025 ACC/AHA/ACEP/NAEMSP/SCAI guideline for the management of patients with acute coronary syndromes: A report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2025 Apr;151(13):e771-e862. doi: 10.1161/CIR.0000000000001309.  
Statin Tolerance
14. Fitchett DH, Hegele RA, Verma S. Statin intolerance. *Circulation* 2015;131:e389-391. <https://doi.org/10.1161/CIRCULATIONAHA.114.013189>
15. Manpuya WM, Cho L, Frid D, et al. Treatment strategies in patients with statin intolerance: the Cleveland Clinic experience. *American Heart Journal* 2013; 166(3):597-603.

16. Zhang H, Plutzky J, Skentzos S, et al. Discontinuation of statins in routine care settings. *Ann of Intern Med* 2013; 158(7):526-534.
17. Backes JM, Ruisinger JF, Gibson CA, et al. Statin-associated muscle symptoms—managing the highly intolerant. *J Clin Lipidol*. 2017 Jan-Feb;11(1):24-33. doi: 10.1016/j.jacl.2017.01.006.
18. Thompson PD, Panza G, Zaleski A, et al. Statin-associated side effects. *J Am Coll Cardiol* 2016 May 24;67(20):2395-2410. doi: 10.1016/j.jacc.2016.02.071.
19. Warden BA, Guyton JR, Kovacs AC, et al. Assessment and management of statin-associated muscle symptoms (SAMS): A clinical perspective from the National Lipid Association. *J Clin Lipidol*. 2023 Jan-Feb;17(1):19-39. doi: 10.1016/j.jacl.2022.09.001.
20. Cheeley MK, Saseen JJ, Agarwala A, et al. NLA scientific statement on statin intolerance: a new definition and key considerations for ASCVD risk reduction in the statin intolerant patient. *J Clin Lipidol*. 2022 Jul-Aug;16(4):361-375. doi: 10.1016/j.jacl.2022.05.068.

**Coding Implications**

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J1306	Injection, inclisiran, 1 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	01.06.22	02.22
Added HCPCS code [J1306].	06.30.22	
Template changes applied to other diagnoses/indications and continued therapy section.	10.06.22	
1Q 2023 annual review: per 2022 ACC expert consensus decision pathway, lowered minimum LDL requirement to 55 mg/dL for members with ASCVD at very high risk and added corresponding Appendix I; references reviewed and updated.	10.18.22	02.23
Per guidelines: for HeFH, added pathway for baseline LDL of at least 160 mg/dL for age < 20 years.	05.17.23	08.23
RT4: added expanded indication to include patients with primary hyperlipidemia per PI. For Commercial LOB, approval duration revised to “6 months or to the member’s renewal date, whichever is longer”.	07.20.23	
1Q 2024 annual review: for redirection to a preferred PCSK9 inhibitor added requirement for 8 week trial duration; added the following requirement from initial approval criteria to also require for continuation of therapy “Treatment plan does not include coadministration with Juxtapid, Repatha, or Praluent”; divided criteria with multiple elements into separate bullets for added	02.09.24	02.24

Reviews, Revisions, and Approvals	Date	P&T Approval Date
clarity; Appendix I clarified smoking is specific to tobacco; references reviewed and updated. Reorganized diagnostic criteria in section I.A for improved clarity (no changes to clinical content).		
1Q 2025 annual review: added contraindication for hypersensitivity per PI; references reviewed and updated.	11.06.24	02.25
RT4: updated indication to reflect revised use as an adjunct to exercise (rather than statin therapy) for hypercholesterolemia per PI; revised hyperlipidemia to hypercholesterolemia throughout the criteria.	08.06.25	

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible

for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note:**

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2022 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.